



## Women's views and postpartum follow-up in the CHIPS Trial (Control of Hypertension in Pregnancy Study)



Marianne Vidler<sup>a</sup>, Laura A. Magee<sup>b,\*</sup>, Peter von Dadelszen<sup>b</sup>, Evelyne Rey<sup>m</sup>, Susan Ross<sup>j</sup>, Elizabeth Asztalos<sup>f,g,n</sup>, Kellie E. Murphy<sup>g,n</sup>, Jennifer Menzies<sup>b</sup>, Johanna Sanchez<sup>n</sup>, Joel Singer<sup>c,r</sup>, Amiram Gafni<sup>d</sup>, Andrée Gruslin<sup>l</sup>, Michael Helewa<sup>i</sup>, Eileen Hutton<sup>k</sup>, Shoo K. Lee<sup>f</sup>, Terry Lee<sup>r</sup>, Alexander G. Logan<sup>h</sup>, Wessel Ganzevoort<sup>o</sup>, Ross Welch<sup>q</sup>, Jim G. Thornton<sup>p</sup>, Jean-Marie Moutquin<sup>e</sup> the CHIPS Study Group<sup>1</sup>

<sup>a</sup>Obstetrics and Gynaecology, University of British Columbia, Canada

<sup>b</sup>Medicine, University of British Columbia, Canada

<sup>c</sup>School of Population and Public Health, University of British Columbia, Canada

<sup>d</sup>Clinical Epidemiology and Biostatistics, McMaster University, Canada

<sup>e</sup>Obstetrics and Gynaecology, Université de Sherbrooke, Canada

<sup>f</sup>Pediatrics, University of Toronto, Canada

<sup>g</sup>Obstetrics and Gynaecology, University of Toronto, Canada

<sup>h</sup>Medicine, University of Toronto, Canada

<sup>i</sup>Obstetrics and Gynaecology, University of Manitoba, Canada

<sup>j</sup>Obstetrics and Gynaecology, University of Alberta, Canada

<sup>k</sup>Obstetrics and Gynaecology, McMaster University, Canada

<sup>l</sup>Obstetrics and Gynaecology, University of Ottawa, Canada

<sup>m</sup>Medicine and Obstetrics and Gynaecology, University of Montreal, Canada

<sup>n</sup>The Centre for Mother, Infant and Child Research, Sunnybrook Research Institute, University of Toronto, Canada

<sup>o</sup>Obstetrics and Gynaecology, University of Amsterdam, Netherlands

<sup>p</sup>Obstetrics and Gynaecology, University of Nottingham, United Kingdom

<sup>q</sup>Obstetrics and Gynaecology, Derriford Hospital, United Kingdom

<sup>r</sup>Centre for Health Evaluation and Outcome Sciences (CHÉOS), Providence Health Care Research Institute, UBC, Vancouver, Canada

### ARTICLE INFO

#### Article history:

Received 4 March 2016

Received in revised form 11 July 2016

Accepted 26 July 2016

#### Keywords:

Hypertension

Blood pressure control

Pregnancy

Postpartum

Satisfaction

### ABSTRACT

**Objective:** To compare women's views about blood pressure (BP) control in CHIPS (Control of Hypertension In Pregnancy Study) (NCT01192412).

**Design:** Quantitative and qualitative analysis of questionnaire responses.

**Setting:** International randomised trial (94 sites, 15 countries).

**Population/sample:** 911 (92.9%) women randomised to 'tight' (target diastolic blood pressure, 85 mmHg) or 'less tight' (target diastolic blood pressure, 100 mmHg) who completed questionnaires.

**Methods:** A questionnaire was administered at ~6–12 weeks postpartum regarding post-discharge morbidity and views about trial participation. Questionnaires were administered by the site coordinator, and contact was made by phone, home or clinic visit; rarely, data was collected from medical records. Quantitative analyses were Chi-square or Fisher's exact test for categorical variables, mixed effects multinomial logistic regression to adjust for confounders, and  $p < 0.001$  for statistical significance. NVivo software was used for thematic analysis of women's views.

**Main outcome measures:** Satisfaction, measured as willingness to have the same treatment in another pregnancy or recommend that treatment to a friend.

**Results:** Among the 533 women in 'tight' ( $N = 265$ ) vs. 'less tight' ( $N = 268$ ) control who provided comments for qualitative analysis, women in 'tight' (vs. 'less tight') control made fewer positive comments about the amount of medication taken (5 vs. 28 women, respectively) and intensity of BP monitoring (7 vs. 17, respectively). However, this did not translate into less willingness to either have the same treatment in another pregnancy (434, 95.8% vs. 423, 92.4%, respectively;  $p = 0.14$ ) or recommend that treatment to a friend (435, 96.0% and 428, 93.4%, respectively;  $p = 0.17$ ). Importantly, although

\* Corresponding author at: Maternal Medicine, St. George's, University of London, Cranmer Terrace, Room J0.27, London SW17 0RE, United Kingdom.

E-mail address: [LMagee@sgul.ac.uk](mailto:LMagee@sgul.ac.uk) (L.A. Magee).

<sup>1</sup> The members of the CHIPS Study Group is listed in Table S1 of [Appendix A](#).

satisfaction remained high among women with an adverse outcome, those in 'tight' control who suffered an adverse outcome (vs. those who did not) were not consistently less satisfied, whereas this was not the case among women in 'less tight' control among whom satisfaction was consistently lower for the CHIPS primary outcome ( $p < 0.001$ ), severe hypertension ( $p \leq 0.01$ ), and pre-eclampsia ( $p < 0.001$ ).

**Conclusions:** Women in 'tight' (vs. 'less tight') control were equally satisfied with their care, and more so in the face of adverse perinatal or maternal outcomes.

© 2016 The Author(s). Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The CHIPS Trial (Control of Hypertension In Pregnancy Study) was an international randomised trial of 'less tight' [target diastolic blood pressure (dBp) of 100 mmHg] vs. 'tight' control (target dBp of 85 mmHg) of non-severe non-proteinuric hypertension in pregnancy [1]. 'Less tight' (vs. 'tight') control was associated with no benefit or harm to the baby, in terms of the primary outcome of perinatal loss or high level neonatal care for >48 h (31.4% vs. 30.7%), or the secondary outcome of serious maternal complications (3.7% vs. 2.0%). However, 'less tight' (vs. 'tight') control was associated with maternal risk, in terms of more severe hypertension (40.6% vs. 27.5%,  $p < 0.0001$ ), platelet count  $<100 \times 10^9/L$  (4.3% vs. 1.6%,  $p < 0.05$ ), and elevated liver enzymes with symptoms (4.3% vs. 1.8%,  $p < 0.05$ ). Although 'less tight' control did not significantly decrease rates of birth weight <10th centile (16.1% vs. 19.8%), the differential rate of 3.7% may be interpreted by some clinicians to be of potential clinical importance.

On balance, 'tight' control of non-severe pre-existing or gestational hypertension in pregnancy appears to be the prudent clinical approach. If a negative impact on fetal growth were true, there was no evidence that this translated into a negative effect on death or illness for the baby, as reflected by the CHIPS primary outcome for which the trial was adequately powered. However, as pregnant women are known to be reluctant to take medication in pregnancy, clinicians may be reluctant to prescribe 'tight'.

After a closed-meeting presentation of the CHIPS results to investigators, more of them indicated that they would recommend 'tight' control (32, 45.7%) rather than 'less tight' control (6, 8.6%), but a significant number (30, 42.9%) were undecided and indicated that they would seek women's views before making their recommendation.

In order to further inform clinicians about 'tight' (vs. 'less tight') control, we extend the findings of the main trial by exploring the views of women who were assigned to each type of BP control and completed a postpartum questionnaire.

## Materials and methods

CHIPS was a pragmatic, open, multicentre, international RCT (ISRCTN 71416914; NCT01192412) [1]. It was approved by the Research Ethics Board at the University of British Columbia as the co-ordinating centre and at all study sites.

The protocol is available on-line at [nejm.org](http://nejm.org). In brief, women at 14<sup>+0</sup>–33<sup>+6</sup> weeks gestation with non-proteinuric pre-existing or gestational hypertension, office diastolic BP (dBp) 90–105 mmHg (or 85–105 mmHg if on antihypertensives) and a live fetus were randomised to 'less tight' control (target dBp 100 mmHg) or 'tight' (target dBp 85 mmHg). The composite primary outcome was pregnancy loss or high level neonatal care for >48 h in the first 28 days of life, and the secondary outcome, serious maternal complications before 6 weeks postpartum. Outcomes were compared between groups using logistic regression adjusted for key prognostic factors.

This report is focussed on information collected by site co-ordinators who were not responsible for direct patient care and

who had health sciences backgrounds and CHIPS-specific training. Co-ordinators administered a structured postpartum questionnaire designed to: (i) collect information about potential post-discharge maternal or neonatal morbidities in the 6 weeks following birth for the mother, or 28 days of life for the baby, whichever was later, that may contribute to the primary or secondary outcomes; and (ii) explore maternal views about trial participation. The questionnaire was completed by contacting all CHIPS participants at 6–12 weeks postpartum (or 36 weeks corrected gestational age for preterm babies, whichever was later), by telephone, home or clinic visit, as appropriate for the centres; if this was not possible, the co-ordinator contacted the woman's medical practitioner and/or reviewed medical records in order to inform the primary and secondary outcomes (but not the satisfaction component of the questionnaire).

Women's views about trial participation were explored by two standard questions: willingness to have the same treatment again in a future pregnancy, and willingness to recommend the same treatment to a friend [2,3]. Women were invited to provide comments after each question, as well as any other additional comments that they wished to share. All responses were recorded by co-ordinators at the time of interview, and subsequently entered verbatim in English into the study database for analysis.

Quantitative analysis of the questionnaire was by Chi-square or Fisher's exact test as appropriate for categorical variables, with  $p < 0.001$  considered to be statistically significant as specified a priori by the CHIPS protocol.

Using mixed effects multinomial logistic regression, we simultaneously estimate the odds ratio (OR) of 'definitely yes' vs. 'probably/definitely not' and 'probably yes' vs. 'probably/definitely not' for 'less tight' vs. 'tight' control. The  $p$  value is against the alternative hypothesis that at least one of these ORs is not equal to 1. In the first analysis as in the primary CHIPS analysis [1], we adjusted for hypertension type, previous severe hypertension, use of any antihypertensive therapy at randomisation, gestational diabetes, weeks of gestation at randomisation, and centre (as random effect). In an additional analysis, we also adjusted for ethnicity, region, and perinatal mortality ratio of the recruiting country to account for potential cultural differences in women's views.

NVivo qualitative research software (Version 10, 2012, QSR International, Doncaster Victoria, Australia) was used to organise, analyse, and interpret women's free text responses. A thematic analysis approach was used for this study. This method is frequently used to identify patterns in large data sets so that relevant patterns can be explored in-depth [4]. Data were coded and stored as nodes that were subsequently used to explore similar and disparate themes that either were specified a priori or emerged during analysis (Panel 1). The first stage of analysis included a 'start list' of codes; these initial themes reflected the expected responses. Subsequently, themes were added, combined and split to better represent the respondents' views that did not reasonably fit into an a priori thematic category. All data were coded by one researcher (MV), a public health researcher with extensive experience in qualitative analyses specifically within the hypertensive disorders of pregnancy. Thematic categories were reviewed by study

Themes defined <i>a priori</i>	Data-driven themes	FINAL THEMES USED
<b>Process of care</b>		
Increased attention from health care providers		Extra health care provider attention
Better quality care		Quality care
Medication/treatment provided*		Quality of treatment
<b>Research processes</b>		
	Comments about research staff	Comments about research staff
Participating in research to help others		Play role in research & help others
	Learning opportunity	Learning opportunity
	Quality of study	Quality of study
Required extra work		Amount of work required
	Importance of research	Importance of research
<b>Drivers of decision-making</b>		
Felt reassured/worried		Feelings of safety (+) or worry (-)
Safety		
Medication/treatment provided*		Amount of medication
Being in control of blood pressure level		Blood pressure level
	Increased BP monitoring	Increased BP monitoring
<b>Outcomes</b>		
Maternal outcomes		Maternal outcomes
Infant outcomes		Infant outcomes
<b>Other</b>		
Any / no problems		Any / no problems
Depends on pregnancy		Planning for future pregnancies
Not applicable		
Don't know		Don't know

\*This theme was divided into 'quality of treatment' and 'amount of medication'.

**Panel 1.** A priori and data driven theme categories for analysis of maternal responses to CHIPS postpartum questionnaire.

investigators (LAM, PvD) who provided input on coding and themes. Respondents were classified to allow for analysis of the qualitative findings according to: allocation to 'less tight' or 'tight' BP control, response to each of the maternal satisfaction questions, and occurrence of each of the following adverse outcomes: primary perinatal, secondary maternal, severe maternal hypertension, pre-eclampsia, delivery at <34 weeks, and delivery at <37 weeks.

## Results

There were 987 women randomised in CHIPS to 'less tight' ( $N = 497$ ) or 'tight' control ( $N = 490$ ) and included in the analyses [1]. For the 981 women with data for the primary and secondary outcomes (493 in 'less tight' and 488 in 'tight' control), an additional 10 (5 vs. 5, respectively) were lost to follow-up for the postpartum questionnaire after delivery, resulting in completed

postpartum questionnaires for maternal and neonatal morbidity for 971 (99.0%) women (Table 1). The primary perinatal outcome (perinatal loss or high level neonatal care for >48 h) was informed by the postpartum questionnaire in 25 (8.2%) babies (15/150 in 'tight' and 10/155 in 'less tight') and the secondary maternal outcome in 5 (17.9%) women (4/10 in 'tight' and 1/18 in 'less tight').

Postpartum questionnaires were completed by mothers at a median of about 8 weeks postpartum. Most questionnaires were completed by telephone interview. In 60 (6.2%) women (30 in 'less tight' and 30 in 'tight' control), questionnaires were completed entirely by use of medical records, 52 because they were not contactable, and 8 because of stillbirth or neonatal death that made the co-ordinators uncomfortable contacting the women (Table S2). As such, there were 911 postpartum questionnaires (from 92.9% of women in the primary analysis) with data informative for the maternal satisfaction and views analyses presented in this paper.

**Table 1**

Postpartum questionnaire completion for the 981 women in CHIPS ( $N$  (%) women or median [IQR], as appropriate).

	'Less tight' control $N = 493$	'Tight' control $N = 488$	$p^a$
Respondents by any method	488 (99.0%)	483 (99.0%)	1.00
Missing	5	5	
Time after delivery questionnaire completed (week)	8.29 [6.57; 11.57]	8.00 [6.71; 10.57]	0.55
Method of questionnaire completion <sup>b</sup>			
Telephone	309 (62.7%)	287 (58.8%)	0.21
Personal interview	158 (32.0%)	166 (34.0%)	0.51
Information obtained (in whole or part) from medical records or maternity care provider	43 (8.7%)	41 (8.4%)	0.86
Missing	5	5	

<sup>a</sup> Using Chi-square.

<sup>b</sup> Responses are not mutually exclusive.

**Table 2**  
Women's views about participation in CHIPS for 911 respondents to maternal satisfaction questions (N (%) women).

	'Less tight' control (N = 458)	'Tight' control (N = 453)	Unadjusted analysis <i>p</i> <sup>a</sup>	Adjusted analyses	
				<i>p</i> <sup>b</sup>	<i>p</i> <sup>c</sup>
Women willing to have the same treatment in another pregnancy					
Respondents (N women)	454 (99.1%)	452 (99.8%)			
<i>Provided no answer</i>	4 (0.9%)	1 (0.2%)			
Definitely yes	309 (67.5%)	307 (67.8%)	0.18	0.15	0.14
Probably yes	114 (24.9%)	127 (28.0%)			
Probably NOT	22 (4.8%)	15 (3.3%)			
Definitely NOT	9 (2.0%)	3 (0.7%)			
Provided further comments for qualitative analysis	268/454 (59.0%)	265/452 (58.6%)			
Would recommend the same treatment to a friend					
Respondents (N women)	455 (99.3%)	453 (100%)			
<i>Provided no answer</i>	3 (0.7%)	0			
Definitely yes	296 (64.6%)	281 (62.0%)	0.27	0.17	0.17
Probably yes	132 (28.8%)	154 (34.0%)			
Probably NOT	24 (5.2%)	16 (3.5%)			
Definitely NOT	3 (0.7%)	2 (0.4%)			
Provided further comments for qualitative analysis	207/455 (45.5%)	213/453 (47.0%)			
Respondents to one/both questions who chose to accept the invitation to provide other comments	241/455 (53.0%)	219/453 (48.3%)			

<sup>a</sup> Using Fisher's exact test.

<sup>b</sup> Adjusted for hypertension type, previous severe hypertension, use of antihypertensive therapy at randomisation (yes/no), gestational diabetes at randomisation, weeks of gestation at randomisation, and centre (as random effect) as in the main CHIPS adjusted analysis [1].

<sup>c</sup> Adjusted for hypertension type, previous severe hypertension, use of antihypertensive therapy at randomisation (yes/no), gestational diabetes at randomisation, weeks of gestation at randomisation, and centre (as random effect) as in the main CHIPS adjusted analysis [1]. In addition to ethnicity, region and perinatal mortality ratio of the recruiting country to account for potential cultural differences in women's views.

### Maternal satisfaction with care

Table 2 presents the women's views about their participation in the CHIPS Trial, from 911 respondents to the maternal satisfaction questionnaire. Just over half of the women (i.e., 268/455 in 'less tight' and 265/453 in 'tight' control) provided additional qualitative comments. The vast majority of women in 'tight' (and 'less tight') control were satisfied with their care, as measured by willingness to either have the same treatment in another pregnancy and highlighted in grey in Table 2 (92.4% in 'less tight' vs. 95.8% in 'tight' control) or recommend the same treatment to a friend (93.4% vs. 96.0%, respectively).

Table 3 presents responses according to trial outcomes for all but the CHIPS secondary outcome for which there were too few events (N = 28) to allow for meaningful interpretation. Overall, more than 80% of women reported being satisfied with their care, regardless of their allocated group or pregnancy outcome; however, women in 'less tight' control with an adverse outcome (vs. those without) appeared to be less satisfied than women in 'tight' control who had an adverse outcome (vs. those without). Among women who received 'less tight' control, those with an adverse outcome (vs. those without) were significantly less satisfied (according to willingness to have the same treatment in a future pregnancy and recommend treatment to a friend) for all outcomes other than birth weight <10th centile. Among women in 'tight' control, women with an adverse outcome (vs. those without) were less willing to have the same treatment in a future pregnancy for the CHIPS primary outcome and pre-eclampsia, but satisfaction measured by recommending treatment to a friend was not significant for any outcome.

### Qualitative analyses according to 'less tight' vs. 'tight' control and satisfaction with care

Following completion of the closed-ended questions; just over half of women provided additional general comments following the first, 'willingness' question, and just under half commented following the second 'recommend to a friend' question (Table 2).

The final list of 18 themes used for qualitative analysis included five themes driven by the data, four in research processes (i.e., comments about research staff, learning opportunity, quality of study, and importance of research) and one in drivers of decision-making (i.e., increased BP monitoring) (Panel 1).

Table 4 presents women's themes according to 'less tight' vs. 'tight' control. Within each treatment group, responses according to whether women were willing to have the same treatment in a future pregnancy (as a measure of satisfaction), and then whether the comments were positive (e.g., worry) or negative (e.g., reassurance) within that theme are presented in Table S3. (Responses for the satisfaction question of 'would recommend treatment to a friend' were similar and are presented in Appendix Table S4.) In 'tight' (and 'less tight') control groups, most themes were commented on positively. Women's statements usually related to processes of care (especially extra health care provider care and quality of care), and less frequently, to 'drivers of decision-making', followed by research-processes and outcomes.

With regards to process of care, comments reflected the importance of one-to-one personalised attention (e.g., "**kept a close eye on me throughout my pregnancy**", 'less tight') and the feeling that the quality of care was better due to participation in the trial (e.g., "**because I seemed to be looked after better**", 'tight'). With regards to the themes related to delivery of care and how the research was conducted, women expressed that the trial enabled them and their families to learn from the experience, as research staff answered questions and explained the condition and procedures thoroughly (e.g., "**everything was explained to me, I understood everything**", 'tight' control). The information led some women to become more involved in their health (e.g., "**made me more involved/compliant with my BP management**", 'tight' control).

With regards to 'drivers of clinical decision-making', women in 'tight' control felt safe (e.g., "**I feel more reassured about my health, my pregnancy, and my baby**", 'tight') and positively about the BP level achieved, as did women in 'less tight' control (e.g., "**my mind was at ease about not taking anything and that I was safe on the trial**", 'less tight'; "**I felt that my blood pressure was well controlled**", 'less tight'). Women in 'tight' (vs. 'less tight') control

**Table 3**  
Satisfaction by outcome among women in 'less tight' (N=488) vs. 'tight' control (N=483).<sup>a</sup>

Theme	'Less Tight' control (N=488)		p	'Tight' control (N=483)		p <sup>b</sup>
	Adverse outcome	No adverse outcome		Adverse outcome	No adverse outcome	
CHIPS primary outcome	N=151	N=337		N=148	N=335	
Willing to have treatment in future pregnancy			<b>0.005*</b>			<b>0.02*</b>
Yes	126 (83.4%)	305 (90.5%)		127 (85.8%)	308 (91.9%)	
No	17 (11.3%)	15 (4.5%)		10 (6.8%)	8 (2.4%)	
Missing	8	17		11	19	
Recommend treatment to a friend			<b>0.002*</b>			0.41
Yes	127 (84.1%)	309 (91.7%)		130 (87.8%)	306 (91.3%)	
No	16 (10.1%)	12(3.6%)		7 (7.5%)	11 (5.4%)	
Missing	8	16		11	18	
Birth weight <10th centile	N=77	N=411		N=95	N=388	
Willing to have treatment in future pregnancy			0.37			0.14
Yes	68 (88.3%)	363 (88.3%)		83 (87.3%)	352 (90.7%)	
No	7 (9.1%)	25 (6.1%)		6 (6.3%)	12 (3.1%)	
Missing	2	23		6	24	
Recommend treatment to a friend			0.43			0.39
Yes	69 (89.6%)	367 (89.3%)		85 (89.5%)	351 (90.5%)	
No	6 (7.8%)	22 (5.3%)		5 (5.3%)	13 (3.4%)	
Missing	2	22		5	24	
Severe hypertension	N=197	N=291		N=131	N=352	
Willing to have treatment in future pregnancy			<b>0.004*</b>			0.08
Yes	170 (86.3%)	261 (89.7%)		113 (86.3%)	322 (91.5%)	
No	21 (10.6%)	11 (3.8%)		8 (6.2%)	10 (2.8%)	
Missing	6	19		10	20	
Recommend treatment to a friend			<b>0.01*</b>			0.93
Yes	173 (87.8%)	263 (90.4%)		117 (89.3%)	319 (90.6%)	
No	18 (9.1%)	10 (3.4%)		5 (3.8%)	13 (3.7%)	
Missing	6	18		9	20	
Pre-eclampsia	N=237	N=251		N=221	N=262	
Willing to have treatment in future pregnancy			<b>0.0005*</b>			<b>0.03*</b>
Yes	200 (84.4%)	231 (92.0%)		193 (87.3%)	242 (92.4%)	
No	25 (10.5%)	7 (2.8%)		13 (5.9%)	5 (1.9%)	
Missing	12	13		15	15	
Recommend treatment to a friend			<b>0.0002*</b>			0.92
Yes	202 (85.2%)	234 (93.2%)		199 (90.0%)	237 (90.5%)	
No	23 (9.7%)	5 (2.0%)		8 (3.6%)	10 (3.8%)	
Missing	12	12		14	15	

<sup>a</sup> Satisfaction was explored according to whether women were willing to have the same treatment in a future pregnancy (yes/no), or recommend the same treatment to a friend in her pregnancy (yes/no).

<sup>b</sup> p value by Chi-square or Fisher's Exact Test, as appropriate.

\* These results were significant with a p-value.

**Table 4**  
Theme categories mentioned by women when asked about willingness to have the same treatment in another pregnancy.<sup>a</sup>

Theme	'Less tight' (N=268)		'Tight' (N=265)	
	Pos	Neg	Pos	Neg
Process of care				
Extra health care provider attention	63	1	60	1
Quality care	60	2	55	0
Quality of treatment	17	5	16	3
Research processes				
Comments about research staff	20	1	21	0
Play role in research & help others <sup>b</sup>	23	0	16	0
Learning opportunity <sup>b</sup>	11	0	26	0
Quality of study	14	1	14	0
Amount of work required	11	4	16	3
Importance of research <sup>b</sup>	2	0	3	0
Drivers of decision-making				
Feelings of safety (+) or worry (-)	23	5	22	3
Amount of medication	28	8	5	9
Blood pressure level achieved	11	14	11	6
Increased BP monitoring	17	1	7	1
Outcomes				
Maternal outcomes	24	10	33	6
Infant outcomes	4	0	13	3
Other				
Any/no problems	14	0	11	0
Planning for future pregnancies <sup>c</sup>	8	0	7	0
Don't know <sup>c</sup>	3	0	6	0

<sup>a</sup> Results are presented according to randomised group ('less tight' vs. 'tight' control), willingness to have the same treatment in another pregnancy ('yes' or 'no'), and whether the comments made were positive or negative.

<sup>b</sup> These themes are positive by their nature.

<sup>c</sup> These themes are neither positive nor negative in nature.

did appear to make fewer positive comments about the amount of medication taken (5 vs. 28, respectively) and the level of BP monitoring (7 vs. 17, respectively), but more positive comments regarding outcomes for both mother (33 vs. 24, respectively) and baby (13 vs. 4, respectively).

Most negative comments were made by women who were not willing to have the same treatment again in a future pregnancy or recommend that treatment to a friend (Table S3). Women were worried about having taken too much medication in 'tight' control (e.g., **"did not like having high doses of medication"**) or too little medication in 'less tight' control (e.g., **"being off meds made her feel anxious and concerned [about] stroke or heart attack"**). Women in 'tight' control may have less frequently expressed concerns about BP level achieved (under drivers of clinical decision-making), and outcomes compared with women in 'less tight' control, as illustrated by comments such as, **"I cannot have less tight control in the future as my last pregnancy was difficult in terms of blood pressure management"** ('less tight'), and **"[I] was not happy with having an abruption, wondering if being in less tight group contributed"** ('less tight').

## Comment

### Summary of results

In CHIPS, more than 90% of women in both 'tight' and 'less tight' control groups expressed satisfaction as reflected by a willingness to have the same treatment in a future pregnancy or recommend that treatment to a friend. Additional comments revealed that women in both BP control groups had a positive research experience and high regard for the quality of care received. This finding likely reflects the high quality of care provided to women as part of the CHIPS Trial.

In 'tight' control, satisfaction remained high among women with an adverse outcome, even in comparison with women without an adverse outcome; this was not consistently true among women in 'less tight' control. Qualitative analysis of comments supported these findings, despite women in 'tight' control making fewer positive comments about the amount of medication received or the level of BP monitoring. This would suggest that preferences to limit medication use and monitoring are not the primary drivers of satisfaction overall or with outcomes. Nevertheless, some women in both groups reported anxieties related to treatment of their BP during pregnancy, highlighting the importance of psychological support in pregnancy and postpartum.

### Strengths and weaknesses

Strengths of this study include its large numbers, randomisation to 'less tight' vs. 'tight' BP control (such that the two groups of women should have differed only by the type of BP control that they received), high response rate, and inclusion of open-ended questions that allowed participants to describe their feelings and explain their responses. Also, the qualitative and quantitative findings are from the same individuals, so the former results can be used to support the latter.

Weaknesses include that the postpartum questionnaire was administered by the CHIPS trial co-ordinators to whom respondents knew from data collection throughout the trial; although women may have felt uncomfortable expressing negative feelings regarding their treatment, the co-ordinators were not in charge of their care. There were small numbers of women in some thematic groups, making inferences difficult. Given the number of countries involved ( $N=15$ ) and languages ( $N=9$ ), it was not possible to either explore responses by region/country of recruitment or conduct the thematic analysis first in the original language which

may have avoided potential error introduced by translation followed by thematic analysis in English. Satisfaction may have been best explored using scales rather than by asking about willingness to have the same treatment again or recommend it to a friend, but CHIPS was a large, international trial, and issues of language, culture and cost influenced our decision to go with simplicity. Postpartum questionnaires were not completed for 8/25 women who suffered perinatal losses; in this minority of cases, study co-ordinators did not feel comfortable contacting participants (as outlined in the CHIPS protocol) to explore satisfaction of the CHIPS Trial. Although we respected their views, and the number of women affected was small, our decision may have skewed the results. Finally, the views presented here may not be generalisable to hypertensive pregnant women in general.

### Comparison with the published literature

Our data from the main CHIPS Trial document similar satisfaction rates to those reported by women who participated in the CHIPS Pilot Trial, in which more than 90% of women were satisfied with their BP management and care, regardless of allocation to 'less tight' or 'tight' BP control [5]. This was true despite being very involved in management of their BP, as in the main CHIPS Trial; women were expected to know the dBp goal to which they had been randomised, and to carry a diary in which all BP measurements were to be recorded. Our data from the main CHIPS Trial go further by collecting and qualitatively analysing additional comments that women were invited to provide. Comments from women assigned to 'tight' (or 'less tight') control most commonly related favourably to process of care. This is consistent with knowledge that women's involvement in their care, the quality of the caregiver-patient relationship, and the amount of support from caregivers all influence women's evaluation of care, regardless of setting [6,7]. We were unable to identify additional published literature on women's views about their BP management per se, in distinction to their preferences for home BP measurement [8], outpatient (vs. inpatient) care [9], and adequate information about prognosis [10]. These studies indicate the preference for outpatient monitoring, similar to the care provided for the majority of women in CHIPS (i.e., 84.0% vs. 83.2% in 'less tight' vs. 'tight' control, respectively) [8,9]. However, numerous studies exploring patient satisfaction in RCTs have reported similar drivers in willingness to participate: personal clinical benefit, a duty to help others, the importance of contributing to research, increased specialist attention, learning, a good rapport with the research team, clinical reassurance, and perceptions of treatment [11–13].

A stakeholder analysis would be ideal to seek the views of clinicians regarding 'tight' and 'less tight' control of BP, taking into account primary and secondary analyses of CHIPS data, including those of women's views.

## Conclusion

This qualitative study extends the findings of the main CHIPS Trial by detailing the individual experiences of women who received 'less tight' and 'tight' control. Women in 'tight' control were satisfied with their care, even in the face of adverse outcomes. Clinicians prescribing 'tight' BP control in pregnancy should feel reassured by this information.

## Funding

The CHIPS Trial was supported by a grant (MCT 87522) from the Canadian Institutes of Health Research.

## Authors' contribution

All named authors contributed to the design of the project, its execution and analysis, and the writing and revision of the manuscript.

## Ethical approval

The CHIPS Trial was approved by the Research Ethics Board at the University of British Columbia as the Co-ordinating Centre (H08-00882) and at all study sites.

## Conflict of interest

Dr. von Dadelszen reports receiving consulting fees and in-kind support of research from Alere International related to preeclampsia and fetal growth restriction through the provision of Triage PLGF cartridges.

Dr. Helewa reports grants from CIHR-Factors associated with inadequate prenatal care, grants from CIHR-Reducing inequities in

access to and use of prenatal care in the Winnipeg Health Region through Health System improvement, grants from CIHR-Quality of prenatal care questionnaire development, grants from CIHR-Effect of Folic Acid supplementation in pregnancy on preeclampsia Folic Acid Clinical Trial (FACT), outside the submitted work. Dr. Magee reports grants from Canadian Institutes of Health Research, during the conduct of the study; personal fees from Bill & Melinda Gates Foundation, outside the submitted work. Jennifer Menzies reports grants, personal fees and non-financial support from CIHR, during the conduct of the study. No other potential conflict of interest relevant to this article was reported.

This manuscript is dedicated to the memory of our dear friend and colleague, Dr. Andrée Gruslin.

## Acknowledgements

We sincerely thank all of the women who gave of themselves by participating in CHIPS. This manuscript is dedicated to the memory of our dear friend and colleague, Dr. Andrée Gruslin.

We thank the 987 women who participated in CHIPS.

## Appendix A

**Table S1**

CHIPS Study Group.

**Steering Committee:** Laura A. Magee (Chair), Elizabeth Asztalos, Amiram Gafni, Andrée Gruslin, Michael Helewa, Eileen Hutton, Shoo Lee, Alexander Logan, Jennifer Menzies, Jean-Marie Moutquin, Kellie Murphy, Evelyne Rey, Sue Ross, Johanna Sanchez, Joel Singer, Peter von Dadelszen

**Working Group:** Laura A. Magee (Chair), Elizabeth Asztalos, Peter von Dadelszen, Trinh Hoac, Joanne Kirton, Jennifer Menzies, Sue Ross, Johanna Sanchez, Katherine Trigiani, Ainy Zahid

**Adjudication Committee:** Laura A. Magee (Chair), Elizabeth Asztalos, Kellie Murphy, Evelyne Rey, Peter von Dadelszen

**Data Safety Monitoring Board:** Michael B. Bracken (Chair), Patricia Crowley, Lelia Duley, Richard Ehrenkranz, Kevin Thorpe

**Data programmers and analysts:** Sunny Chan, Michael Shi, Shelley Yu

**Collaborators:** The number of women recruited in each country and centre is specified in brackets.

**ARGENTINA (36):**

**Hospital LC Lagomaggiore,** Mendoza (16): Raquel de Lourdes Martin, Maria Florencia Bassi, Mirta Clara Caruso, Valeria Lagunas, Fernando Vera

**Hospital Avellaneda,** Tucuman (10): Maria Mohedano de Duhalde, Alicia Beatriz Roque, Patricia Roldan, Esteban Marcos Duhalde, Viviana Dip

**Hospital JR Vidal,** Corrientes (8): Jesus Daniel Aguirre, Elba Mirta Alicia Morales, Griselda Itati Abreo, Teresa De Sagastizabal, Carolina Gomez, Nadia Rizzi

**Hospital JM Cullen,** Santa Fe (2): Carlos Arias, Ricardo Antonio Bruno

**AUSTRALIA (85):**

**Ipswich Hospital,** Ipswich (36): Kassam Mahomed, Alison Drew, Ann Green, Jane Hoare

**Women's and Children's Hospital,** Adelaide (18): Bill Hague, Suzette Coat, Caroline Crowther, Peter Muller, Sophie Trenowden

**King Edward Memorial Hospital,** Subiaco (17): Barry Walters, Claire Parker, Dorothy Graham, Craig Pennell, Eileen Sung

**Campbelltown Hospital,** Penrith South (8): Angela Makris, Gaksoo Lee, Charlene Thornton, Annemarie Hennessy

**Liverpool Hospital,** Penrith South (5): Angela Makris, Gaksoo Lee, Charlene Thornton, Annemarie Hennessy

**St John of God Hospital,** Subiaco (1): Louise Farrell, Claire Parker, Eileen Sung, Barry Walters

**BRAZIL (19):**

**Maternidade de Vila Nova Choeirinha,** Sao Paulo (7): Nelson Sass, Henri Korkes, Dayana Couto Ferreira

**Hospital Universitario Antonio Pedro,** Niteroi (6): Renato Augusto Moreira de Sa, Monique Schmidt Marques Abreu

**Maternidade Escola da UFRJ,** Rio de Janeiro (4): Rita Guerios Bornia, Nancy Ribeiro da Silva, Fernanda Freitas Oliveira Cardoso

**Hospital Sao Lucas – PUCRS,** Porto Alegre (2): Caio Coelho Marques, Jorge Hornos, Ricardo Leal Davdt, Letícia Germany Paula, Pedro Luis Zanella

**CANADA (233):**

**British Columbia Women's Hospital and Health Centre,** Vancouver (32): Laura A. Magee, Peter von Dadelszen, Gabrielle Inglis, Ruth Dillon, Ashley Docherty, Anna Hutfield

**Jim Pattison Outpatient Care and Surgery Centre,** Surrey (26): Keith Still, Sayrin Lalji, Tamara Van Tent, Chris Hotz, Tracy Messmer

**St Michael's Hospital,** Toronto (22): Joel G. Ray, Howard Berger, Leanne De Souza, Andrea Lausman, Tatiana Freire-Lizama, Kate Besel

**Foothills Medical Centre,** Calgary (21): Paul Gibson, Greta Ellsworth, Leslie Miller, T. Lee-Ann Hawkins

**Sunnybrook Health Sciences Centre,** Toronto (19): Michelle Hladunewich, Anna Rogowsky, Dini Hui, Virginia Collins

**IWK Health Centre,** Halifax (19): Isabelle Delisle, Cora Fanning

**Royal Alexandra Hospital,** Edmonton (16): Nestor Demianczuk, Rshmi Khurana, Winnie Sia, Catherine Marnoch, Carmen Young, Cheryl Lux

**CHU Sainte-Justine,** Montreal (15): Evelyne Rey, Sophie Perreault, Valerie Tremblay

**CHUS Fleurimont,** Sherbrooke (13): Anne-Marie Côté, Jean-Marie Moutquin, Sophie Desindes, Veronique Dagenais

**Ottawa Hospital Civic Division,** Ottawa (13): Andrée Gruslin, Heather Clark, Elaine O'Shea, Ruth Rennicks White

**Mount Sinai Hospital,** Toronto (8): Shital Gandhi, Mary-Jean Martin, Cheryl Brush, Gareth Seaward

**Royal University Hospital,** Saskatoon (6): Jill Newstead-Angel, Judy Brandt, Jocelyne Martel, Kristine Mytopher, Elise Buschau

**Ottawa Hospital General Division,** Ottawa (5): Andree Gruslin, Erin Keely, Patti Waddell, Ruth Rennicks White, Svetlana Shachkina, Alan Karovitch

**St Paul's Hospital,** Vancouver (5): Robert Anderson, Nicole Koenig, Theresa Yong

**Toronto East General Hospital,** Toronto (5): Marie Vasiliou, Peri Johnson, Beth Allan

**London Health Sciences Centre,** London (4): Renato Natale, Laura Kennedy

**Royal Victoria Hospital,** Montreal (2): Lucie Opatrny, Lorraine Lavigne

**Regina General Hospital,** Regina (1): George Carson, Sheila Kelly

**Women's Health Centre,** St John's (1): Joan Crane, Donna Hutchens

**CHILE (57):**

**Hospital Dr Sotero del Rio,** Puente Alto (45): Juan Pedro Kusanovic, Christian Figueroa, Karla Silva Neculman, Juan Andres Ortiz, Paula Vargas

(Continued)

- Hospital Base Osorno**, Osorno (12): Pedro Ferrand, Jorge Carrillo
- COLOMBIA** (36):  
**Corporacion Comfenalco Valle - Universidad Libre** (20), **Clinica Versalles** (11), **Clinica Materno Infantil Farallones** (5), Cali: Rodrigo Cifuentes Borrero, Dahiana Marcela Gallo, Luisa Fernanda Moreno
- ESTONIA** (19):  
**Tartu University Hospital - Women's Clinic**, Tartu (19): Fred Kirss, Kristiina Rull, Anne Kirss
- HUNGARY** (5):  
**University of Debrecen**, Debrecen (5): Tamas Major, Andrea Fodor, Tunde Bartha
- ISRAEL** (12):  
**Hillel Yaffe Medical Center**, Hadera (6): Mordechai Hallak, Nardin Aslih, Saja Anabousi-Murra, Ester Pri-Or  
**Ma'ayney Hayeshua Medical Center**, Bnei Brak (3): Linda Harel, Sima Siev  
**Nazareth Hospital (EMMS)**, Nazareth (3): Marwan Hakim, Christina Simona Khoury, Najla Hamati
- JORDAN** (13):  
**Islamic Hospital**, Amman (13): Mazen El-Zibdeh, Lama Yousef
- NEW ZEALAND** (17):  
**Christchurch Women's Hospital**, Christchurch (16): Ruth Hughes, Di Leishman, Barbra Pullar  
**Waitemata Health-North Shore Hospital**, Auckland (1): Matthew Farrant
- POLAND** (21):  
**Medical University of Gdansk**, Gdansk (9): Malgorzata Swiatkowska-Freund, Krzysztof Preis, Anette Aleksandra Traczyk-Los, Anna Partyka, Joanna Preis-Orlikowska, Mariusz Lukaszuk  
**Polish Mothers Memorial Hospital**, Lodz (9): Grzegorz Krasomski, Michaela Krekora, Anna Kedzierska-Markowicz, Katarzyna Zych-Krekora  
**University School of Medical Sciences**, Poznan (3): Grzegorz H. Breborowicz, Anna Dera-Szymanowska
- THE NETHERLANDS** (96):  
**Academic Medical Center**, Amsterdam (28): Wessel Ganzevoort, Jannet Bakker, Joost Akkermans, Anouk Pels  
**OLVG**, Amsterdam (13): Eline van den Akker, Sabine Logtenberg  
**UMCU**, Utrecht (10): Steven Koenen, Maartje de Reus, David Borman, Martijn A. Oudijk  
**VU Medical Center**, Amsterdam (9): Annemieke Bolte, Viki Verfaillie, Bart Graaf  
**Maxima Medical Centre**, Veldhoven (8): Martina Porath, Corine Verhoeven, Ben Willem Mol  
**UMCG**, Groningen (6): Maureen T.M. Franssen, Lida Ulkeman, Ineke Hamming, Jose H.M. Keurentjes, Ina van der Wal  
**Isala Klinieken Zwolle**, Zwolle (5): S.W.A. Nij Bijvank, A.A. Lutjes  
**Tergoozieziekenhuizen**, Hilversum (5): Henricus Visser, Jannet Bakker  
**MUMC Maastricht**, Maastricht (4): Hubertina Catharina Johanna Scheepers  
**St Antonius Ziekenhuis**, Nieuwegein (3): Erik van Beek, David Borman, Coby van Dam, Kathy van den Berg-Swart  
**Kennemer Gasthuis Haarlem**, Haarlem (2): Paula Pernet, Birgit van der Goes  
**Diakonessen Ziekenhuis**, Utrecht (1): Nico Schuitemaker  
**Flevo ziekenhuis**, Almere (1): Gunilla Kleiverda, Marcel van Alphen, Ageeth Rosman  
**Jeroen Bosch Hospital**, 's-Hertogenbosch (1): Ingrid Gaugler-Senden, Marieke Linders
- UNITED KINGDOM** (268):  
**Guy's & St Thomas' Hospital**, London (38): Catherine Nelson-Piercy, Annette Briley, May Ching Soh, Kate Harding, Hayley Tarft  
**New Cross Hospital**, Wolverhampton (31): David Churchill, Katherine Cheshire, Julia Icke, Mausumi Ghosh  
**Nottingham City Hospital**, Nottingham (30): James Thornton, Yvonne Toomassi, Karen Barker, Joanne Fisher, Nicky Grace, Amanda Green, Joanne Gower, Anna Molnar, Shobhana Parameshwaran, Andrew Simm  
**Queen's Medical Centre**, Nottingham (22): James Thornton, George Bugg, Yvette Davis, Ruta Desphande, Yvette Gunn, Mohammed Houda, Anna Molnar, Nia Jones  
**Royal Victoria Infirmary**, Newcastle upon Tyne (22): Jason Waugh, Carly Allan, Gareth Waring  
**Liverpool Women's Hospital**, Liverpool (16): Steve A. Walkinshaw, Angela Pascall, Mark Clement-Jones, Michelle Dower, Gillian Houghton, Heather Longworth, Tej Purewal  
**Bradford Royal Infirmary**, Bradford (13): Derek Tuffnell, Diane Farrar, Jennifer Syson, Gillian Butterfield, Vicky Jones, Rebecca Paethorpe, Tracey Germaine  
**Leicester Royal Infirmary**, Leicester (12): Marwan Habiba, Debbie Lee  
**Wexham Park Hospital**, Slough (12): Olufemi Eniola, Lynne Blake, Jane Khan  
**City Hospitals Sunderland NHS Foundation Trust**, Sunderland (10): Helen M. Cameron, Kim Hinshaw, Amanda Bargh, Eileen Walton  
**South Warwickshire NHS Trust**, Warwick (9): Olanrewaju Sorinola, Anna Guy, Zoe D'Souza, Rhiannon Gabriel, Jo Williams  
**Derriford Hospital**, Plymouth (8): Ross Welch, Heidi Hollands  
**York Hospital**, York (8): Olujimi Jibodu, Sara Collier, Pauline Tottie, Claire Oxby, James Dwyer  
**Singleton Hospital**, Swansea (7): Franz Majoko, Helen Goldring, Sharon Jones  
**Chesterfield Royal Hospital**, Chesterfield (6): **Janet Cresswell**, Louise Underwood, Mary Kelly-Baxter, Rebecca Robinson  
**Sheffield Teaching Hospitals NHS Foundation Trust**, Sheffield (6): Dilly Anumba, Anne Chamberlain, Clare Pye  
**St Mary's Hospital**, Manchester (6): **Clare Tower**, Sue Woods, Lisa Horrocks, Fiona Prichard, Lynsey Moorhead, Sarah Lee, Louise Stephens, Cara Taylor, Suzanne Thomas, Melissa Whitworth, Jenny Myers  
**Birmingham Women's Hospital**, Birmingham (5): Ellen Knox, Katie Freitas, Mark Kilby, Amanda Cotterill  
**Lancashire Teaching Hospitals NHS Foundation Trust**, Lancashire (3): Khalil Abdo, Katrina Rigby, Julie Butler, Fiona Crosfill, Sean Hughes, Sanjeev Prashar, Fatimah Soydemir  
**The Royal Derby Hospital**, Derby (3): Janet Ashworth, Lorraine Mycock, Jill Smith  
**Basildon & Thurrock University Hospital**, Basildon (1): **Amaju Ikomi**, Kerry Goodsell, Jean Byrne, Maxwell Masuku, Alice Pilcher
- USA** (70):  
**Cooper University Hospital**, Camden (13): Meena Khandelwal, Gunda Simpkins, Michelle Iavicoli, Yon Sook Kim, Richard Fischer, Robin Perry  
**Medical University of South Carolina**, Charleston (11): Eugene Y. Chang, Tamara D. Saunders, Betty W. Oswald, Kristin D. Zaks  
**Beth Israel Deaconess**, Boston (8): Sarosh Rana, Dawn McCullough  
**Yale-New Haven Hospital**, New Haven (8): Anna Sfakianaki, Cheryl Danton, Erin Kustan, Luisa Coraluzzi  
**Norton Hospital Downtown** (7), **Norton Suburban Hospital** (2), Louisville: Helen How, Christina Waldon  
**East Carolina University**, Greenville (6): Jeffrey Livingston, Sherry Jackson, Lisa Greene  
**Meriter Hospital**, Madison (6): Dinesh Shah  
**Oregon Health & Science University**, Portland (5): Jorge E. Tolosa, Monica Rincon, Leonardo Pereira, Amy E. Lawrence, Janice E. Snyder  
**University of North Carolina**, Chapel Hill (4): D. Michael Armstrong, Teresa Blue, Austin Hester, Kathryn Salisbury

## Appendix B. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ejogrb.2016.07.509>.

## References

- [1] Magee LA, von DP, Rey E, et al. Less-tight versus tight control of hypertension in pregnancy. *N Engl J Med* 2015;372(5):407–17.
- [2] Junttila EK, Karjalainen PK, Ohtonen PP, Raudaskoski TH, Ranta PO. A comparison of paracervical block with single-shot spinal for labour analgesia in multiparous women: a randomised controlled trial. *Int J Obstet Anesth* 2009;18(1):15–21.
- [3] Santos AR, Bahamondes MV, Hidalgo MM, Atti A, Bahamondes L, Monteiro I. Pain at insertion of the levonorgestrel-releasing intrauterine system in nulligravida and parous women with and without cesarean section. *Contraception* 2013;88(1):164–8.
- [4] Braun V, Clarke V. Thematic analysis. Washington: American Psychological Association; 2012.
- [5] Magee LA, von Dadelszen P, Chan S, et al. The Control of Hypertension in Pregnancy Study Pilot Trial. *BJOG* 2007;114(6):770. e13–770, e20.
- [6] Hodnett ED. Pain and women's satisfaction with the experience of childbirth: a systematic review. *Am J Obstet Gynecol* 2002;186(5 Suppl. Nature):S160–72.
- [7] Srivastava A, Avan BI, Rajbangshi P, Bhattacharyya S. Determinants of women's satisfaction with maternal health care: a review of literature from developing countries. *BMC Pregnancy Childbirth* 2015;15(1):97.
- [8] Taylor RS, Freeman L, North RA. Evaluation of ambulatory and self-initiated blood pressure monitors by pregnant and postpartum women. *Hypertens Pregnancy* 2001;20(1):25–33.
- [9] Turnbull DA, Wilkinson C, Gerard K, et al. Clinical, psychosocial, and economic effects of antenatal day care for three medical complications of pregnancy: a randomised controlled trial of 395 women. *Lancet* 2004;363(9415):1104–9.
- [10] Leeners B, Rath W, Kuse S, Neises M, Neumaier-Wagner P. Satisfaction with medical information in women with hypertensive disorders in pregnancy. *J Psychosom Res* 2006;60(1):39–44.
- [11] Canvin K, Jacoby A. Duty, desire or indifference? A qualitative study of patient decisions about recruitment to an epilepsy treatment trial. *Trials* 2006;7:32.
- [12] Locock L, Smith L. Personal experiences of taking part in clinical trials – a qualitative study. *Patient Educ Couns* 2011;84(3):303–9.
- [13] McCann SK, Campbell MK, Entwistle VA. Reasons for participating in randomised controlled trials: conditional altruism and considerations for self. *Trials* 2010;11:31.