

Distribution of bacteriophages in food and environment samples

^{2,3*}Premarathne, J. M. K. J. K., ²Thung, T. Y., ²New, C. Y., ⁴Huat, J. T. Y., ⁵Basri, D. F.,
²Rukayadi, Y., ⁶Nakaguchi, Y., ⁷Nishibuchi, M. and ^{1,2}Son, R.

¹Food Safety and Food Integrity, Institute of Tropical Agriculture and Food Security, Universiti Putra Malaysia 43400 Serdang, Selangor, Malaysia

²Department of Food Science, Faculty of Food Science and Technology, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

³Department of Livestock and Avian Science, Faculty of Livestock, Fisheries and Nutrition, Wayamba University of Sri Lanka, Makandura, 60170 Gonawila, Sri Lanka

⁴Faculty of Food Technology, Universiti Sultan Zainal Abidin, 20400 Kuala Terengganu, Terengganu, Malaysia

⁵School of Diagnostic and Applied Health Sciences, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 UKM Kuala Lumpur, Selangor Darul Ehsan, Malaysia

⁶Faculty of Bioresources and Environmental Sciences, Ishikawa Prefectural University, Nonoichi City, Ishikawa Prefecture, Japan

⁷Centre for Southeast Asian Studies, Kyoto University, Kyoto 606-8501, Japan

Article history

Received: 9 November 2016
 Received in revised form:
 20 December 2016
 Accepted: 2 January 2017

Abstract

Foodborne pathogens have become a constant threat to the consumer and food industry. Reduce efficacy of antibiotics with emergence of resistant bacteria has limited the opportunities for controlling pathogenic bacteria in food commodities and treating foodborne infections. Bacteriophages can be a promising alternative for alleviate the risk of transmitting pathogenic bacteria via food commodities. Therefore, this research was conducted to find distribution of bacteriophages in diverse niches in order to identify suitable sources for isolating bacteriophages to use controlling foodborne pathogens. Firstly bacterial strains were screened for lysogenic and selected suitable host bacterial strains were used for isolating and determining bacteriophage titer in fresh raw food and environmental samples. Eighteen different lytic bacteriophages effective against *Campylobacter*, *S. aureus*, *L. monocytogenes* and *E. coli* were isolated from this study. Bacteriophages titer was determined within range of 10^2 to 10^{10} PFU/mL and bacteriophages were most frequently isolated from chicken (60%) samples. The isolated bacteriophages could be potential candidates for controlling foodborne diseases.

Keywords

Foodborne pathogens
 Bacteriophages
 Double layer assay
 Plaques
 Bacteriophage titer

© All Rights Reserved

Introduction

Campylobacter, *Staphylococcus aureus*, *Listeria monocytogenes* and *Escherichia coli* are some of the leading foodborne pathogens that cause major public health issue around the world (Behravesh *et al.*, 2012; Newell *et al.*, 2010). Food products, especially food of animal origin can be contaminated with these foodborne pathogens at any stage during the production process (Sillankorva *et al.*, 2012; Carvalho *et al.*, 2010).

High *Campylobacter* contamination level was reported in chicken and vegetables sold in Malaysia (Tang *et al.*, 2009; Chai *et al.*, 2007; Wong *et al.*, 2017). Dairy and milk products sold in Malaysian markets were contaminated with *S. aureus* (Sasidharan *et al.*, 2011). Neela *et al.* (2009) reported the isolation of MRSA from pigs and pig handlers in

Malaysia. *L. monocytogenes* recovered from various food sources such as ready to eat foods (Marian *et al.*, 2012), ducks (Adzitey *et al.*, 2013), raw salad vegetables (Ponniah *et al.*, 2010), burger patties (Wong *et al.*, 2012), chicken (Goh *et al.*, 2013) and chicken offal (Kuan *et al.*, 2013). The *E. coli* O157:H7 occurred in beef samples (Son *et al.*, 1998); milk (Lye *et al.*, 2013); ready-to-eat food, popiah (Elexson *et al.*, 2017) and chicken (Chang *et al.*, 2013) sold in Malaysia. Also, *E. coli* was isolated from raw vegetables (Loo *et al.*, 2013), pigs (Ho *et al.*, 2013) and water samples (Alhaj *et al.*, 2007) in the country. Also high antibiotic resistance was reported among the isolated foodborne pathogens (Tang *et al.*, 2009; Chai *et al.*, 2007). Though not significant, GMO that may carry the antibiotic resistant gene as marker has also been reported in Malaysia (Lisha *et al.*, 2017).

Therefore, to combat foodborne illnesses

*Corresponding author.
 Email: krissjyaruk@yahoo.com

and antibiotic resistance, novel, and efficient strategies have become paramount in the current global context (Gálvez *et al.*, 2010; García *et al.*, 2010). Bacteriophages have been identified as a potential alternative to antibiotics (Haq *et al.*, 2012; Sillankorva *et al.*, 2012). Bacteriophages also known as phages are bacterial viruses composed of nucleic acid genome enclosed within a protein or lipoprotein coat (Sillankorva *et al.*, 2012). Phages are ubiquitous in the environment and all most all the bacteria possess their own bacteriophages that can be isolated from the environment where the host bacteria occur (Sillankorva *et al.*, 2012; Kutter, 2005).

More than 1031 bacteriophages contain in the earth biosphere (Endersen *et al.*, 2014). Further, bacteriophages that infect various host bacteria were isolated from diverse sources such as human faeces, animal faeces, food, water, soil and sewage (Atterbury, 2009). However, worldwide still an only limited number of bacteriophages were identified (Casjens, 2008). Even in Malaysia, only a few bacteriophages such as that effective against *Vibrio cholera* (Al-Fendi *et al.*, 2014) and colibacillosis (Lau *et al.*, 2010) were investigated. Therefore, need to study and isolate prevailing bacteriophages in the country that could be used as a potential agent to control pathogenic and multidrug resistance bacteria. Therefore, the objectives of this study were to assess the distribution and concentration of *Campylobacter*, *S. aureus*, *L. monocytogenes*, and *E. coli* bacteriophages in food and environmental samples.

Materials and Methods

Host-bacterial cultures and growth conditions

Foodborne pathogenic bacteria including *Campylobacter jejuni*, *Campylobacter coli* isolated from this study and *Staphylococcus aureus*, methicillin-resistant *Staphylococcus aureus* (MRSA), *Listeria monocytogenes* and *E. coli* obtained from the Food Safety Laboratory, Faculty of Food Science and Technology, Universiti Putra Malaysia were used as the target host bacteria.

Campylobacter strains used in this study were maintained in 20% (v/v) glycerol supplemented with Brain Heart Infusion (BHI) broth (Merck, Germany) at -20°C until use. When necessary, working cultures were prepared on Blood Agar plates (PB0114, Oxoid, UK). *Campylobacter* host strains were grown according to the method by Carvalho *et al.* (2010). Briefly, *Campylobacter* was grown in New Zealand Casamino Yeast Medium (NZCYM, Sigma-Aldrich, USA) supplemented with 400 $\mu\text{g}/\text{mL}$ CaCl_2 and 400 $\mu\text{g}/\text{mL}$ MgSO_4 at 42°C under

microaerophilic conditions generated by Anaerocult C system (Merck, Germany) with shaking (150 rpm) (MaxQ 4000; Barnstead International, USA). The broth reached the mid-log phase were used for bacteriophage enrichment.

The bacterial strains, *S. aureus*, *L. monocytogenes* and *E. coli* utilized in this study, were stored in 20% (v/v) glycerol at -20°C . A 100 μL of the frozen stock culture was inoculated into Luria Bertani (LB) broth (Merck, Germany) and incubated overnight at 37°C with shaking. Secondary cultures were prepared on Tryptone Soy agar (TSA) (Merck, Germany) slants by transferring from the activated bacterial broth culture grown in Luria-Bertani (LB) broth (Merck, Germany). Then the agar slants were sealed and stored at 4°C until use for the preparation of working cultures. When necessary a loop full of culture from agar slants was transferred onto TSA (Merck, Germany) (incubated overnight at 37°C), then checked for purity and transferred single colony into LB broth (Merck, Germany) to prepare working cultures. The bacterial host strains (*S. aureus*, *L. monocytogenes*, and *E. coli*) grown in LB broth (Merck, Germany) with shaking incubation (150 rpm) at 37°C until it reaches the mid-log phase were used for bacteriophage enrichment.

Screening bacterial strains for lysogeny

Bacterial strains were screened for lysogeny by using mitomycin C as inducing agent. A 10 mL of fresh NZCYM broth growth media was inoculated with 100 μL of an overnight culture of the *Campylobacter* strain to be tested for lysogeny. The *Campylobacter* cultures were incubated at 42°C under microaerobic conditions (Anaerocult C system (Merck, Germany)). Once the bacteria reached the exponential growth phase, 5 μL of 1 mg/mL mitomycin C was added and for the negative control 5 μL of distilled water was added. Then absorbency was measured at 600 nm in each hour for 12 h. The strains that reduced turbidity were excluded from the study.

A 10 mL of fresh LB broth growth media was inoculated with 100 μL of an overnight culture of the *S. aureus*, *L. monocytogenes* or *E. coli* strain to be tested for lysogeny. Then cultures were incubated at 37°C , and once they reached the exponential growth phase, 5 μL of 1 mg/mL mitomycin C was added and for the negative control 5 μL of distilled water was added. Then absorbency was measured at 600 nm in each hour for 8 h. The strains that reduced turbidity were excluded from the study.

Sample collection

Fresh food samples including; dairy products,

poultry, seafood, vegetables, and beef were purchased from different wet markets in Selangor, Malaysia. Also, environmental samples such as water and sewage samples were collected from surrounding area of the wet markets in Selangor and Universiti Putra Malaysia, Malaysia. Samples were collected and stored inside an insulated cool box containing ice pack and immediately processes at the laboratory.

Enrichment of bacteriophages

Samples were mixed with 1:10 (w/v) SM buffer [50 mM Tris-HCl (pH 7.5)], 0.1 M NaCl, 8 mM MgSO₄•7H₂O and 0.01% (w/v) gelatine (Sigma-Aldrich, USA) and the mixture was pumped. Then the homogenised sample was inoculated with exponential phase *Campylobacter* culture at 9:1 (v/v) ratio. Then followed by overnight shaking incubation (150 rpm) at 42°C microaerobically (Anaerocult C system; Merck, Germany) for potential bacteriophages to amplify and dissociate. The homogenized sample was inoculated with exponential phase bacterial culture of *S. aureus*, MRSA, *L. monocytogenes*, and *E. coli* at 9:1 (v/v) ratio and followed by overnight shaking incubation (150 rpm) at 37°C for potential bacteriophages to amplify and dissociate.

Isolation of bacteriophages

After incubation, the mixture was centrifuged for 10 min at 10,000 rpm to remove bulk debris from the sample. Bacterial cells were removed by filtering the supernatant through 0.2 µm membrane filter (Sartorius, Germany). Then, the filtrate was checked for phage activity using spot plate assay (Chang *et al.*, 2005; Adam, 1959). Briefly, an aliquot of 100 µL of mid-log phase bacteria *Campylobacter* cultures were inoculated into 3 mL of molten NZCYM soft agar (NZCYM broth containing 0.6% (w/v) agar). This mixture was poured evenly onto NZCYM bottom agar (NZCYM broth containing 1.5% (w/v) agar). Once the top agar solidified, 10 µL of the filtrate was spotted onto the onto NZCYM bottom agar containing *Campylobacter*. Then the plates were dried for 30 mins and incubated overnight at 42°C under microaerobic conditions (Anaerocult C system; Merck, Germany).

An aliquot of 100 µL of mid-log phase bacterial cultures of *S. aureus*, MRSA, *L. monocytogenes* and *E. coli* were separately inoculated into 3 mL of molten LB soft agar (LB broth containing 0.6% (w/v) agar). This mixture was poured evenly onto LB agar (Merck, Germany). Once the top agar solidified, 10 µL of the filtrate was spotted onto the LB agar containing host bacteria (*S. aureus*, MRSA, *L. monocytogenes*, and *E. coli*). Then the plates were dried for 30 mins and

all culture plates were incubated overnight at 37°C.

Plaque morphology

After incubation, the plates were observed for the presence of visible clear zones or plaques that are indicative of bacterial lysis by phage activity. Plaques were scored on a scale from 0 to 5 depending on the plaque morphology (Al-Fendi, 2014). Plates with no interaction or minimal effect were given a score 0 and 1 respectively. A Clear and complete bacterial lysis was scored as 2 while turbid plaques that were not able to clearly distinguish as lytic activity was given a score 3 or 4. Plaque with a clear margin and turbid centre that resembles a bulls-eye was given score 5. Initially, plaques scored 2-5 were considered as phage activity.

Double layer assay

Samples that displayed phage activity were reconfirmed using double layer assay as described by Adams (1959). Briefly, a 100 µL of enriched sample filtrate and 100 µL mid-log phase *Campylobacter* bacterial culture was added to 3 mL of molten NZCYM soft agar (NZCYM broth containing 0.6% (w/v) agar). Then the mixture was mixed well and poured evenly onto NZCYM bottom agar (NZCYM broth containing 1.5% (w/v) agar). Once the top agar solidified, the plates were incubated overnight at 42°C under microaerobic conditions (Anaerocult C system; Merck, Germany).

Briefly, a 100 µL of enriched sample filtrate and 100 µL mid-log phase bacterial culture of *S. aureus*, MRSA, *L. monocytogenes*, and *E. coli* was separately added to 3 mL of molten LB soft agar (LB broth containing 0.6% (w/v) agar). The mixture was mixed well and poured evenly onto LB agar (Merck, Germany). Once the top agar solidified, the culture plates were incubated overnight at 37°C.

After incubation, plates were observed for the presence of plaques. Samples indicated presences of plaques were confirmed as phages. Detected plaques were picked using a sterile pipette tip and resuspended in 100-500 µL of SM buffer and stored at 4°C for future use.

Bacteriophages titration

Bacteriophages titration was conducted using the double layer assay as described by Adams (1959). A ten folds dilution of phage lysate was prepared by using SM buffer. Then double layer assay was conducted for each dilution in triplicates.

After incubation, the plates were observed for the presence of visible clear zones of bacterial lysis. The dilution that formed 30 and 200 plaques were selected

and counted the number of plaques. The obtained data were used to calculate the plaque forming units (PFU/ mL) using standard formula (Adams, 1959).

Results and Discussion

Screening for lysogeny

Out of the 36 tested bacterial strains for prophage activity using mitomycin C, two of the *S. aureus* strains harboured prophages in the genome. Incubating with mitomycin C, the optical density reduced in the bacterial strains that harboured prophage. At the initial phase of phage isolation, selecting bacterial hosts that do not harbour prophages in the genome is important (Garcia *et al.*, 2007). The prophages in the bacterial genome can be induced by application of mitomycin C, chloroform or ultraviolet (UV) light (Weinbauer, 2004; Garcia *et al.*, 2007). In this study, mitomycin C was used to induce the prophages, and two *S. aureus* strains (FQ3 and FQ5) were identified to be lysogenize by containing prophages in the genome. Therefore, those two *S. aureus* strains excluded from phage isolation procedure.

Plaque formation

First samples were screened using the spot plate assay, observed various plaque morphologies were given a score and recorded. Different scores given for plaque morphologies indicate in Figure 1. Initially, 131 experiments were able to produce clear to turbid plaques from the spot assay method. In the spot plate method, once the enriched sample applied to a lawn of host bacteria on agar medium, phages in the sample will be absorbed to the host bacterium and commence lysis. This bacterial lysis produces a visible clear zone on the bacterial lawn that named as a plaque (Adams, 1959; Kutter, 2005; Yoon *et al.*, 2007). Samples demonstrated bacteriophage activities on spot plate method was then reconfirmed with double layer assay. In the food samples tested, highest phage activity was demonstrated in seafood samples however with confirmation by double layer assay identified as lysogenic phages (data not shown). However, following the double layer assay, only 18 samples were able to produce clear plaques characteristic of a lytic phage while the rest formed turbid, very small or hardly visible plaques typical for lysogenic phages (Adams, 1959; Yoon *et al.*, 2007). The temperate phages lead to lysogenic infection in which transfers the phage genome into the bacterial progeny without lysis of the host bacterium. Phages that enter lysogenic growth cycle produce turbid plaques on susceptible bacterial lawns (Adams, 1959; Ai *et al.*, 2008). The lytic nature of the phages to be




Figure 1. Observed plaque morphologies

verified before using for phage therapy because often lysogenic phages may not be likely to inhibit bacterial pathogens and might become resistant to phage lysis (Skurnik *et al.*, 2007; Sulakvelidze, 2011). The isolated phages produced around 1.0 to 3.0 mm lytic plaques in the double layer assay (Figure 2). The size and appearance of a plaque can be associated with the volume and density of agar, concentration and stage of growth of the host bacterium and constancy of the top agar (Adams, 1959; Cormier and Janes, 2014).

Occurrence of phages

A total of 18 bacteriophages confirmed to demonstrate lytic activity for *L. monocytogenes*, *E. coli*, *S. aureus*, MRSA, and *Campylobacter* (Table 1). From the tested samples, the highest frequency of isolating bacteriophages was reported in chicken (60%) whilst vegetables found to have the lowest frequency (9.4%) (Table 1). The occurrence of bacteriophages was closely related to the host bacterium (Akhtar *et al.*, 2014). Therefore, bacteriophages could be isolated from natural environment of the targeted host bacterium (Adams, 1959; Endersen *et al.*, 2014). However, none of the phage activity showed by environmental samples was able to produce lytic activity (Table 1). Similarly, Bigwood and Hudson, (2009) were unable to isolate *Campylobacter* phage from water samples taken from various locations can be related to a low concentration of host bacterium

Table 1. Isolation of phages from different samples

Sample Type	Number of <i>L. monocytogenes</i>		<i>E. coli</i> (%)	<i>S. aureus</i> (%)	MRSA (%)	<i>C. jejuni</i> (%)	<i>C. coli</i> (%)	Total (%)
	samples tested	(%)						
Beef	15	1 (6.7)	1 (6.7)	1 (6.7)	1 (6.7)	0 (0.0)	0 (0.0)	4 (26.7)
Chicken	15	4 (26.7)	1 (6.7)	1 (6.7)	0 (0.0)	3 (20)	1 (6.7)	9 (60)
Vegetables	32	1 (3.1)	1 (3.1)	0 (0.0)	0 (0.0)	2 (6.3)	0 (0.0)	3 (9.4)
Clam	10	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Cookies	10	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Shrimp	10	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Environmental samples	20	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	112	6 (5.4)	3 (2.7)	2 (1.8)	1 (0.9)	5 (4.5)	1 (0.9)	18 (16.1)




Figure 2. Morphology of lytic plaques. Phage active against A: *L. monocytogenes*; B: *S. aureus*; C: *C. jejuni*; D: *E. coli*.

present in the environmental samples (Bigwood and Hudson, 2009).

In this study, six different phages effective for *Campylobacter* were isolated from chicken and vegetable samples. Similar to these findings, previously phages effective against *Campylobacter* were isolated from variety of samples including; chicken (Atterbury *et al.*, 2003), chicken intestine (Carrillo *et al.*, 2007; Hansen *et al.*, 2007; Carvalho *et al.*, 2010), duck intestine (Hansen *et al.*, 2007), chicken ceca (Atterbury *et al.*, 2005; El-Shibiny *et al.*, 2005; Carrillo *et al.*, 2007), chicken feces (Owens *et al.*, 2013), and from abattoir wastewater (Hansen *et al.*, 2007). However, phages for *Campylobacter* could not be isolated from water samples taken from rivers, streams, ponds and lakes (Bigwood and Hudson, 2009) and frozen chicken (Atterbury *et al.*, 2003). Interestingly, this study was able to isolate two phages effective against *C. jejuni* from vegetable

samples. High level of *Campylobacter* contamination in vegetables was previously reported from Malaysia (Chai *et al.*, 2007). Phages are naturally present in the environment in which the host bacterium inhabits (Brüssow, 2002). Therefore, an abundance of *Campylobacter* host in vegetables may have associated with the isolation of phages acting against *C. jejuni*.

In the current study, the bacteriophages effective against *E. coli* were isolated each from chicken (n=1), beef (n=1) and vegetables (n=1). Previously *E. coli* phages were isolated from; sewage sludge (Fan *et al.*, 2012), human stools (Tomat *et al.*, 2013), chicken meat (Shousha *et al.*, 2015), poultry feces (Bhensdadia *et al.*, 2014) and industrial cucumber fermentation (Lu and Breidt, 2015).

Three bacteriophages effective against *S. aureus* were isolated from chicken and beef samples and one of which was effective against MRSA. Previously, phages effective against *S. aureus* were isolated from a cow infected with mastitis (Kwiatek *et al.*, 2012), sewage samples (Synnott *et al.*, 2009; Alves *et al.*, 2014) and endotracheal tubes used by patients (Hsieh *et al.*, 2011). The isolated phage ØMRSA1 lysed both *S. aureus* and MRSA strains.

Listeria phages (5.4%) were the most frequently isolated phages in this study, a total of six phages from chicken (n=4), beef (n=1) and vegetables (n=1) that effective against *L. monocytogenes*. Earlier *Listeria* phages were recovered from fish waste treatment seafood (Arachchi *et al.*, 2013), turkey processing plant (Kim *et al.*, 2008), sewage of a dairy processing facility (Carlton *et al.*, 2005) and environmental samples (Loessner and Busse, 1990). Though more than 500 different phages effective for *Listeria* were isolated; only a few were studied at the genomic level (Zimmer *et al.*, 2003; Carlton *et al.*, 2005; Dorscht *et al.*, 2009; Schmuki *et al.*, 2012).

Table 2. Bacteriophage Titers

Source	Bacterial Host	Bacteriophage	Bacteriophage titer (PFU/mL)
Chicken	<i>L. monocytogenes</i>	ØLM1	$5.3 \times 10^3 \pm 3.21$
Chicken	<i>L. monocytogenes</i>	ØLM2	$5.0 \times 10^4 \pm 2.14$
Chicken	<i>L. monocytogenes</i>	ØLM3	$1.3 \times 10^7 \pm 5.26$
Chicken	<i>L. monocytogenes</i>	ØLM6	$2.0 \times 10^7 \pm 3.44$
Beef	<i>L. monocytogenes</i>	ØLM4	$7.0 \times 10^8 \pm 2.51$
Vegetables	<i>L. monocytogenes</i>	ØLM5	$4.7 \times 10^2 \pm 1.04$
Beef	<i>E. coli</i>	ØEC1	$1.78 \times 10^{10} \pm 2.23$
Chicken	<i>E. coli</i>	ØEC2	$8.1 \times 10^{10} \pm 4.02$
Vegetables	<i>E. coli</i>	ØEC3	$4.6 \times 10^3 \pm 1.21$
Beef	<i>S. aureus</i>	ØSA1	$3.0 \times 10^3 \pm 2.11$
Vegetables	<i>S. aureus</i>	ØSA2	$1.1 \times 10^3 \pm 3.47$
Beef	MRSA	ØMRSA1	$4.4 \times 10^4 \pm 3.31$
Chicken	<i>C. coli</i>	ØCC1	$7.27 \times 10^{10} \pm 2.15$
Chicken	<i>C. jejuni</i>	ØCJ1	$9.73 \times 10^{10} \pm 3.64$
Chicken	<i>C. jejuni</i>	ØCJ2	$8.78 \times 10^{10} \pm 2.37$
Chicken	<i>C. jejuni</i>	ØCJ3	$9.90 \times 10^8 \pm 2.56$
Vegetables	<i>C. jejuni</i>	ØCJ4	$3.87 \times 10^6 \pm 2.04$
Vegetables	<i>C. jejuni</i>	ØCJ5	$8.40 \times 10^4 \pm 2.11$

Bacteriophage titers

The bacteriophage titer levels in samples ranged from $8.1 \times 10^{10} \pm 4.02$ to $4.7 \times 10^2 \pm 1.04$ (Table 2). The highest bacteriophage titer at 1010 PFU/mL was detected for *Campylobacter* and *E. coli* specific bacteriophages that isolated from the chicken and beef. While the lowest titer was recorded for phage isolated from vegetables that active against *L. monocytogenes*. Comparatively vegetable samples carried low phages titer than meat samples. Atterbury et al., (2003) was able to isolate 1×10^2 to 4×10^6 PFU/mL *Campylobacter* phages from chicken and later Atterbury et al., (2005) and El-Shibiny et al., (2005) recovered 10^2 to 10^7 PFU/g *Campylobacter* phages from the cecal content of broiler chickens. Enrichment incorporated in the bacteriophage isolation step was reported to facilitated isolation of phages (Carvalho et al., 2010) may be associated with the favourable titers reported in this study.

Conclusion

In conclusion, phages specific for *Campylobacter jejuni*, *C. coli*, *Staphylococcus aureus*, *MRSA*, *Listeria monocytogenes*, and *Escherichia coli* were isolated from various food commodities in Malaysia. Therefore, findings of this study indicate that phages can be readily isolated from food products, and

people can be exposed to phages frequently via food. The high titers and lytic ability exhibited by these phages stipulate them as promising and potential candidates to use as effective biocontrol agents against pathogenic bacteria.

Acknowledgements

This research was supported by the Fundamental Research Grant Scheme (FRGS) of Ministry of Higher Education (MOHE), Malaysia with project code 02-01-14-1475FR and reference code FRGS/1/2014/SG05/UPM/01/2 and in part, by a fund for Research on international cooperation in medical science, Research on global health issues, Health and Labor Science Research Grants, the Ministry of Health, Labor, and Welfare of Japan, and a fund from Kyoto University Research Coordination Alliance, Japan, by a fund by Kakenhi Grant-in-Aid for Scientific Research and from the Japan Society for the Promotion of Sciences.

References

- Adams, M. H. 1959. Appendix: methods of study of bacterial viruses. Bacteriophages, p. 450-451. New York: Interscience Publisher.
- Adzitey, F., Ali, G. R. R., Huda, N., Cogan, T. and Corry, J. 2013. Prevalence, antibiotic resistance and genetic diversity of *Listeria monocytogenes* isolated from

- ducks, their rearing and processing environments in Penang, Malaysia. *Food Control* 32(2): 607-614.
- Ai, Y., Meng, F. and Zeng, Y. 2000. The evolution of pathogen-host interactions mediated by bacteriophages. Wei Sheng Wu Hsueh Pao – *Acta Microbiologica Sinica* 40(6): 657-660.
- Al-Fendi, A. 2014. Isolation and molecular characterization of bacteriophages from environmental sources: Toward the development of phage-based therapy for *Vibrio cholerae*. Penang, Malaysia: The Universiti Sains Malaysia, PhD thesis.
- Al-Fendi, A., Shueb, R. H., Ravichandran, M. and Yean, C. Y. 2014. Isolation and characterization of lytic vibriophage against *Vibrio cholerae* O1 from environmental water samples in Kelantan, Malaysia. *Journal of Basic Microbiology* 54(10): 1036-1043.
- Alhaj, N., Mariana, N. S., Raha, A. R. and Ishak, Z. 2007. Prevalence of antibiotic resistance among *Escherichia coli* from different sources in Malaysia. *International Journal of Poultry Science* 6(4): 293-297.
- Alves, D. R., Gaudion, A., Bean, J. E., Esteban, P. P., Arnot, T. C., Harper, D. R., Kot, W., Hansen, L. H., Enright, M.C. and Jenkins, A. T. A. 2014. Combined use of bacteriophage K and a novel bacteriophage to reduce *Staphylococcus aureus* biofilm formation. *Applied and Environmental Microbiology* 80(21): 6694-6703.
- Arachchi, G. J. G., Mutukumira, A. N., Dias-Wanigasekera, B. M., Cruz, C. D., McIntyre, L., Young, J., Flint, S. H., Hudson, A. and Billington, C. 2013. Characteristics of three listeriaphages isolated from New Zealand seafood environments. *Journal of Applied Microbiology* 115:1427-1438.
- Atterbury, R. J. 2009. Bacteriophage biocontrol in animals and meat products. *Microbial Biotechnology* 2(6): 601-612.
- Atterbury, R. J., Connerton, P. L., Dodd, C. E., Rees, C. E. and Connerton, I. F. 2003. Isolation and characterization of *Campylobacter* bacteriophages from retail poultry. *Applied and Environmental Microbiology* 69(8): 4511-4518.
- Atterbury, R. J., Dillon, E., Swift, C., Connerton, P. L., Frost, J.A., Dodd, C. E. R., Rees, C. E. D. and Connerton, I. F. 2005. Correlation of *Campylobacter* bacteriophage with reduced presence of hosts in broiler chicken ceca. *Applied and Environmental Microbiology* 71(8): 4885-4887.
- Behraves, C. B., Williams, I. T. and Tauxe, R. V. 2012. Emerging foodborne pathogens and problems: expanding prevention efforts before slaughter or harvest. In Choffnes, E. R., Relman, D. A., Olsen, L., Hutton, R., and Mack, A. (Eds). *Improving food safety through a one health approach: workshop summary*. p. 307-330. National Academies Press.
- Bhensdadia, D. V., Bhimani, H. D., Nathani, N.M., Rawal, C. M., Koringa, P. G., Joshi, C.G., Kothari, C.R. and Kothari, R. K. 2014. Isolation, Molecular Characterization and Insight into the Genome Sequence of *E. coli* Bacteriophage ADB-2 from Poultry Fecal Sample. *Journal of Next Generation: Sequencing and Applications* 1(1): 2-7.
- Bigwood, T. and Hudson, J. A. 2009. *Campylobacters* and bacteriophages in the surface waters of Canterbury (New Zealand). *Letters in Applied Microbiology* 48(3): 343-348.
- Brüssow, H. and Hendrix, R. W. 2002. Phage genomics: small is beautiful. *Cell* 108(1): 13-16.
- Canchaya, C., Fournous, G., Chibani-Chennoufi, S., Dillmann, M. L. and Brüssow, H. 2003. Phage as agents of lateral gene transfer. *Current Opinion in Microbiology* 6(4): 417-424.
- Carlton, R. M., Noordman, W. H., Biswas, B., De Meester, E. D. and Loessner, M. J. 2005. Bacteriophage P100 for control of *Listeria monocytogenes* in foods: genome sequence, bioinformatic analyses, oral toxicity study, and application. *Regulatory Toxicology and Pharmacology* 43: 301-312.
- Carrillo, C. M. L., Connerton, P. L., Pearson, T. and Connerton, I. F. 2007. Free-range layer chickens as a source of *Campylobacter* bacteriophage. *Antonie Van Leeuwenhoek* 92(3): 275-284.
- Carvalho, C., Susano, M., Fernandes, E., Santos, S., Gannon, B., Nicolau, A., Gibbs, P., Teixeira, P. and Azeredo, J. 2010. Method for bacteriophage isolation against target *Campylobacter* strains. *Letters in Applied Microbiology* 50(2): 192-197.
- Casjens, S. R. 2008. Diversity among the tailed-bacteriophages that infect the Enterobacteriaceae. *Research in Microbiology* 159(5): 340-348.
- Chai, L. C., Robin, T., Ragavan, U. M., Gunsalam, J. W., Bakar, F. A., Ghazali, F. M., Son, R. and Kumar, M. P., 2007. Thermophilic *Campylobacter* spp. in salad vegetables in Malaysia. *International Journal of Food Microbiology* 117(1): 106-111.
- Chang, H. C., Chen, C. R., Lin, J. W., Shen, G. H., Chang, K. M., Tseng, Y. H. and Weng, S. F. 2005. Isolation and characterization of novel giant *Stenotrophomonas maltophilia* phage ϕ SMA5. *Applied and Environmental Microbiology* 71(3): 1387-1393.
- Chang, W. S., Afsah Hejri, L., Rukayadi, Y., Khatib, A., Lye, Y. L., Loo, Y. Y., Mohd Shahril, N., Soopna, P., Kuan, C. H., Goh, S. G., Tang, J. Y. H. and Son, R. 2013. Quantification of *Escherichia coli* O157: H7 in organic vegetables and chickens. *International Food Research Journal* 20(2): 1023-1029.
- Cormier, J. and Janes, M. 2014. A double layer plaque assay using spread plate technique for enumeration of bacteriophage MS2. *Journal of Virological Methods* 196: 86-92.
- Daubin, V. and Ochman, H. 2004. Bacterial genomes as new gene homes: the genealogy of ORFans in *Escherichia coli*. *Genome Research* 14(6): 1036-1042.
- Dobrindt, U. and Reidl, J. 2000. Pathogenicity islands and phage conversion: evolutionary aspects of bacterial pathogenesis. *International Journal of Medical Microbiology* 290(6): 519-527.
- Dorscht, J., Klumpp, J., Biemann, R., Schmelcher, M., Born, Y., Zimmer, M., Calendar, R. and Loessner, M. J. 2009. Comparative genome analysis of *Listeria* bacteriophages reveals extensive mosaicism, programmed translational frameshifting, and a novel

- prophage insertion site. *Journal of Bacteriology* 191(23): 7206-7215.
- Elexson, N., Nik Yuhanis, F.N., Malcolm, T.T.H., New, C.Y., Chang, W.S., Ubong, A., Kuan, C.H., Loo, Y.Y., Thung, T.Y. and Son, R. 2017. Occurrence of *Escherichia coli* harbouring stx genes in popiah, a Malaysian street food. *Food Research* 1(1): 29-32.
- El-Shibiny, A., Connerton, P. L. and Connerton, I. F. 2005. Enumeration and diversity of campylobacters and bacteriophages isolated during the rearing cycles of free-range and organic chickens. *Applied and Environmental Microbiology* 71(3): 1259-1266.
- Endersen, L., O'Mahony, J., Hill, C., Ross, R. P., McAuliffe, O. and Coffey, A. 2014. Phage therapy in the food industry. *Annual Review of Food Science and Technology* 5: 327-349.
- Fan, H., Fan, H., Mi, Z., Fan, J., Zhang, L., Hua, Y., Wang, L., Cui, X., Zhang, W., Zhang, B., Huang, Y. and Li, J., 2012. A fast method for large-scale isolation of phages from hospital sewage using clinical drug-resistant *Escherichia coli*. *African Journal of Biotechnology* 11(22): 6143-6148.
- Gálvez, A., Abriouel, H., Benomar, N. and Lucas, R. 2010. Microbial antagonists to food-borne pathogens and biocontrol. *Current Opinion in Biotechnology* 21: 142-148.
- García, P., Madera, C., Martínez, B. and Rodríguez, A. 2007. Biocontrol of *Staphylococcus aureus* in curd manufacturing processes using bacteriophages. *International Dairy Journal* 17(10): 1232-1239.
- García, P., Rodríguez, L., Rodríguez, A. and Martínez, B. 2010. Food biopreservation: promising strategies using bacteriocins, bacteriophages and endolysins. *Trends in Food Science and Technology* 21: 373-382.
- Goh, S. G., Kuan, C. H., Loo, Y. Y., Chang, W. S., Lye, Y. L., Soopna, P., Tang, J. Y. H., Nakaguchi, Y., Nishibuchi, M., Afsah-Hejri, L. and Son, R. 2012. *Listeria monocytogenes* in retailed raw chicken meat in Malaysia. *Poultry science* 91(10): 2686-2690.
- Hansen, V. M., Rosenquist, H., Baggesen, D. L., Brown, S. and Christensen, B. B. 2007. Characterization of *Campylobacter* phages including analysis of host range by selected *Campylobacter* Penner serotypes. *BMC Microbiology* 7(1): 90.
- Haq, I. U., Chaudhry, W. N., Akhtar, M. N., Andleeb, S. and Qadri, I. 2012. Bacteriophages and their implications on future biotechnology: a review. *Virology Journal* 9(1): 1-8.
- Helms, M., Simonsen, J., Olsen, K. E. and Mølbak, K. 2005. Adverse health events associated with antimicrobial drug resistance in *Campylobacter* species: a registry-based cohort study. *Journal of Infectious Diseases* 191(7): 1050-1055.
- Ho, W. S., Tan, L. K., Ooi, P. T., Yeo, C. C. and Thong, K. L. 2013. Prevalence and characterization of verotoxigenic-*Escherichia coli* isolates from pigs in Malaysia. *BMC Veterinary Research* 9(1): 109.
- Hsieh, S. E., Lo, H. H., Chen, S. T., Lee, M. C. and Tseng, Y. H. 2011. Wide host range and strong lytic activity of *Staphylococcus aureus* lytic phage Stau2. *Applied and Environmental Microbiology* 77(3): 756-761.
- Kim, J. W., Siletzky, R. M. and Kathariou, S. 2008. Host ranges of *Listeria*-specific bacteriophages from the turkey processing plant environment in the United States. *Applied and Environmental Microbiology* 74: 6623-6630.
- Kuan, C. H., Kuan, C. H., Goh, S. G., Loo, Y. Y., Chang, W. S., Lye, Y. L., Puspanadan, S., Tang, J. Y. H., Nakaguchi, Y., Nishibuchi, M., Mahyudin, N. A. and Son, R. 2013. Prevalence and quantification of *Listeria monocytogenes* in chicken offal at the retail level in Malaysia. *Poultry science* 92(6): 1664-1669.
- Kutter, E. S. A. 2005. *Bacteriophages Biology and Applications*. CRC Press.
- Kwiatek, M., Parasion, S., Mizak, L., Gryko, R., Bartoszcze, M. and Kocik, J. 2012. Characterization of a bacteriophage, isolated from a cow with mastitis, that is lytic against *Staphylococcus aureus* strains. *Archives of Virology* 157(2): 225-234.
- Lau, G. L., Sieo, C. C., Tan, W. S., Hair-Bejo, M., Jalila, A. and Ho, Y. W. 2010. Efficacy of a bacteriophage isolated from chickens as a therapeutic agent for colibacillosis in broiler chickens. *Poultry Science* 89(12): 2589-2596.
- Lisha, V., New, C.Y., Nishibuchi, M. and Son, R. 2017. Rapid genetically modified organism (GMO) screening of various food products and animal feeds using multiplex polymerase chain reaction (PCR). *Food Research* 1: 1-6.
- Loessner, M. J. and M. Busse. 1990. Bacteriophage typing of *Listeria* species. *Applied and Environmental Microbiology* 56(6): 1912-1918.
- Loo, Y. Y., Soopna, P., Goh, S.G., Kuan, C. H., Chang, W. S., Lye, Y. L., Tang, J. Y. H., Rukayadi, Y., Nakaguchi, Y., Nishibuchi, M. and Son, R.. 2013. Quantitative detection and characterization of Shiga toxin-producing *Escherichia coli* O157 and non-O157 in raw vegetables by MPN-PCR in Malaysia. *International Food Research Journal* 20(6): 3313-3317.
- Lu, Z. and Breidt, F. 2015. *Escherichia coli* O157:H7 bacteriophage Φ241 isolated from industrial cucumber fermentation at high acidity and salinity. *Frontiers in Microbiology* 6: 1-10.
- Lye, Y. L., Son, R., Afsah Hejri, L., Chang, W. S., Loo, Y. Y., Soopna, P., Kuan, C. H., Mohd Shahril, N., Rukayadi, Y., Khatib, A., Tang, J.Y.H. and Nishibuchi, M. 2013. Risk of *Escherichia coli* O157:H7 transmission linked to the consumption of raw milk. *International Food Research Journal* 20(2): 1001-1005.
- Marian, M. N., Aminah, S.S., Zuraini, M. I., Son, R., Maimunah, M., Lee, H. Y., Wong, W.C. and Elexson, N. 2012. MPN-PCR detection and antimicrobial resistance of *Listeria monocytogenes* isolated from raw and ready-to-eat foods in Malaysia. *Food Control* 28(2): 309-314.
- Neela, V., Zafrul, A. M., Mariana, N. S., Van Belkum, A., Liew, Y. K. and Rad, E. G. 2009. Prevalence of ST9 methicillin-resistant *Staphylococcus aureus* among pigs and pig handlers in Malaysia. *Journal of Clinical Microbiology* 47(12): 4138-4140.

- Newell, D. G., Koopmans, M., Verhoef, L., Duizer, E., Aidara-Kane, A., Sprong, H., Opsteegh, M., Langelaar, M., Threlfall, J., Scheutz, F. and van der Giessen, J. 2010. Food-borne diseases—the challenges of 20 years ago still persist while new ones continue to emerge. *International Journal of Food Microbiology* 139: S3-S15.
- Owens, J., Barton, M. D. and Heuzenroeder, M. W. 2013. The isolation and characterization of *Campylobacter jejuni* bacteriophages from free range and indoor poultry. *Veterinary Microbiology* 162(1): 144-150.
- Ponniah, J., Robin, T., Paie, M.S., Son, R., Ghazali, F. M., Kqueen, C. Y., Nishibuchi, M., Nakaguchi, Y. and Malakar, P. K. 2010. *Listeria monocytogenes* in raw salad vegetables sold at retail level in Malaysia. *Food Control* 21(5): 774-778.
- Sasidharan, S., Prema, B. and Latha, L. Y. 2011. Antimicrobial drug resistance of *Staphylococcus aureus* in dairy products. *Asian Pacific Journal of Tropical Biomedicine* 1(2): 130-132.
- Schmuki, M. M., Erne, D., Loessner, M. J. and Klumpp, J. 2012. Bacteriophage P70: unique morphology and unrelatedness to other *Listeria* bacteriophages. *Journal of Virology* 86: 13099–13102.
- Shousha, A., Awaiwanont, N., Sofka, D., Smulders, F.J., Paulsen, P., Szostak, M.P., Humphrey, T. and Hilbert, F. 2015. Bacteriophages isolated from chicken meat and the horizontal transfer of antimicrobial resistance genes. *Applied and Environmental Microbiology* 81(14): 4600-4606.
- Sillankorva, S. M., Oliveira, H. and Azeredo, J. 2012. Bacteriophages and their role in food safety. *International Journal of Microbiology* ID 863945: 1-13.
- Skurnik, M., Pajunen, M. and Kiljunen, S. 2007. Biotechnological challenges of phage therapy. *Biotechnology Letters* 29(7): 995-1003.
- Son, R., Mutalib, S.A., Rusul, G., Ahmad, Z., Morigaki, T., Asai, N., Kim, Y.B., Okuda, J. and Nishibuchi, M. 1998. Detection of *Escherichia coli* O157: H7 in the beef marketed in Malaysia. *Applied and Environmental Microbiology* 64(3): 1153-1156.
- Sulakvelidze, A. 2011. The challenges of bacteriophage therapy. *European Industrial Pharmacy* 10: 14-18.
- Synnott, A. J., Kuang, Y., Kurimoto, M., Yamamichi, K., Iwano, H. and Tanji, Y. 2009. Isolation from sewage influent and characterization of novel *Staphylococcus aureus* bacteriophages with wide host ranges and potent lytic capabilities. *Applied and Environmental Microbiology* 75(13): 4483-4490.
- Tang, J. Y. H., Mohamad Ghazali, F., Abdul Aziz, S., Nishibuchi, M. and Son, R. 2009. Comparison of thermophilic *Campylobacter* spp. occurrence in two types of retail chicken samples. *International Food Research Journal* 16(3): 277-288.
- Tomat, D., Mercanti, D., Balagué, C. and Quiberoni, A. 2013. Phage biocontrol of enteropathogenic and Shiga toxin-producing *Escherichia coli* during milk fermentation. *Letters in Applied Microbiology* 57(1): 3-10.
- Weinbauer M. 2004. Ecology of procaryotic viruses. *FEMS Microbiology Reviews* 28: 127–181.
- Wong, C.Y., Usha, M., Vivian, C.Y., Ubong, A., Nakaguchi, Y., Nishibuchi, M. and Son, R. 2017. Cross contamination of *Campylobacter jejuni* from naturally contaminated chicken liver and chicken legs. *Food Research* 1(2): 33-37.
- Wong, W. C., Pui, C.F., Robin, T., Anyi, U., Sahroni, M., Hidayah, N., Mohamad Ghazali, F., Ahmad, N., Cheah, Y. K. and Son, R. 2012. Antibioqram pattern among cultures of *Listeria monocytogenes* isolated from frozen burger patties in Malaysia. *Pertanika Journal of Tropical Agricultural Science* 35(4): 793-804.
- Yoon S. S., Barrangou-Pouey R., Breidt F. and Fleming H. P. 2007. Detection and characterization of a lytic *Pediococcus* bacteriophage from the fermenting cucumber brine. *Journal of Microbiology and Biotechnology* 17(2): 262–270.
- Zimmer, M., Sattelberger, E., Inman, R., Scherer, S., Calendar, R. and Loessner, M. J. 2003. Genome and proteome of *Listeria monocytogenes* bacteriophage PSA: an unusual case for programmed+1 translational frameshifting in structural protein synthesis. *Molecular Microbiology* 50: 303–317.